



**USAID**  
FROM THE AMERICAN PEOPLE

HEALTH POLICY  
INITIATIVE



# **COSTING MALE CIRCUMCISION IN LESOTHO, SWAZILAND, AND ZAMBIA: IMPLICATIONS FOR THE COST- EFFECTIVENESS OF CIRCUMCISION AS AN HIV INTERVENTION**

**SEPTEMBER 2007**

This publication was produced for review by the U.S. Agency for International Development (USAID). It was prepared by staff of the USAID | Health Policy Initiative, Task Order I.

The USAID | Health Policy Initiative, Task Order I, is funded by the U.S. Agency for International Development under Contract No. GPO-I-01-05-00040-00, beginning September 30, 2005. Task Order I is implemented by Constella Futures, in collaboration with the Centre for Development and Population Activities (CEDPA), White Ribbon Alliance for Safe Motherhood (WRA), and Religions for Peace.

# **COSTING MALE CIRCUMCISION IN LESOTHO, SWAZILAND, AND ZAMBIA: IMPLICATIONS FOR THE COST- EFFECTIVENESS OF CIRCUMCISION AS AN HIV INTERVENTION**

**SEPTEMBER 2007**

The views expressed in this publication do not necessarily reflect the views of the U.S. Agency for International Development or the U.S. Government.

# TABLE OF CONTENTS

<b>Acknowledgments</b> .....	<b>iv</b>
<b>Executive Summary</b> .....	<b>v</b>
<b>Introduction</b> .....	<b>1</b>
<b>Methodology</b> .....	<b>1</b>
Cost Analysis .....	1
<b>Unit Cost Estimation</b> .....	<b>5</b>
Direct and Indirect Costs .....	9
Complications and Related Costs .....	11
<b>Scaling Up Circumcision</b> .....	<b>13</b>
Impact and Cost-effectiveness of Scaling up MC.....	13
Pace of Scale-up.....	14
Multiplicative Impact of MC .....	16
Implications to the Health System .....	18
Behavioral Responses to MC.....	19
Sustainability.....	20
Limitations .....	20
<b>Conclusion</b> .....	<b>20</b>
<b>Appendix A. Epidemiology Modeling</b> .....	<b>22</b>
<b>Appendix B. Data Collection Instrument</b> .....	<b>23</b>
<b>Appendix C. Examples of Indirect Costs</b> .....	<b>34</b>
<b>Appendix D. Assumptions Used for Unit Cost Estimation of Comprehensive Package     of Services</b> .....	<b>35</b>
<b>Appendix E. Equipment Requirements</b> .....	<b>37</b>

## ACKNOWLEDGMENTS

The authors (Gayle Martin,<sup>1</sup> Lori Bollinger,<sup>2</sup> Tanvi Pandit-Rajani,<sup>1</sup> Relebohile Tshehlo,<sup>3</sup> R. Nkambula,<sup>4</sup> Steven Forsythe,<sup>2</sup> and John Stover<sup>2</sup>) gratefully acknowledge the Lesotho and Swaziland ministries of health and social welfare (MoHSWs) for their leadership and direction; the clinical and administrative staff of various facilities for their data inputs; the national drug supply organizations (NDSOs) for their assistance with identifying the costs of supplies and drugs; the MoHSWs' Department of Human Resources for providing staffing costs and related expenditures; and the members of the Male Circumcision Task Force for their advice and suggestions on methodology. The authors also acknowledge the Zambia Ministry of Health for its collaboration on this project, as well as Richard Hughes and Evans Chinkoyo of JHPIEGO for their advice and suggestions on methodology.

---

<sup>1</sup> Constella Futures, USAID | Health Policy Initiative, Task Order 1, Washington, DC.

<sup>2</sup> Futures Institute, Glastonbury, CT.

<sup>3</sup> Ministry of Health and Social Welfare, Maseru, Lesotho.

<sup>4</sup> Ministry of Health, Mbabane, Swaziland.

# EXECUTIVE SUMMARY

## Background

Clinical trials have now confirmed the efficacy of male circumcision (MC) in reducing female-to-male HIV transmission. Some cost data have been reported (ranging between US\$25 and US\$69), and these cost data also formed the basis of a cost-effectiveness analysis. It is unclear, however, what exactly is included in the costing studies and hence whether these costs are directly comparable. For example, often, indirect costs are not fully reflected; donations (especially clinicians' time) are not costed; and variation by provider type and level of health facility is not considered. It is anticipated that this cost analysis will provide a more detailed examination of the costs of MC and inform a sounder basis for an assessment of the cost-effectiveness of MC and planning for the implementation of MC. The analysis has three components: (1) conducting key informant interviews to better understand the social, cultural, and policy context of MC; (2) costing MC; and (3) modeling the impact of MC on the HIV epidemic. The qualitative component was only carried out in Lesotho and Swaziland, and results from these interviews can be found in the individual country reports.

## Design/Methodology

### Cost Analysis

The cost analysis is divided into two components: (1) defining the intervention (reviewing existing literature and protocols and conducting key informant interviews with current and potential providers of circumcision); and (2) costing adult male circumcision (collecting direct and indirect cost data—financial, human resources, drugs, supplies, and equipment costs—from providers). An ingredients approach to costing was followed, whereby all the inputs were listed and their contribution to the overall cost was then quantified. Multiple countries and providers (private, government, nongovernmental, missionary) were considered, allowing for standardization, comparison, and validation. The unit costs were adjusted for the probability of and cost associated with complications. Although generally viewed as part of a comprehensive MC package of services,<sup>5</sup> some MC services were not routinely implemented (e.g., pre- and post-circumcision behavioral counseling and HIV testing and training). The costs of these activities were taken from existing service programs that might be unrelated to MC, such as counseling and testing.

### Epidemiologic Modeling

We estimated the impact of male circumcision on the number of new HIV infections using a computer simulation model, Spectrum. The model replicated the dynamics of the HIV epidemic in Lesotho, Swaziland, and Zambia by dividing the population into various risk groups: those not sexually active, those with a single sexual partner, those with more than one sexual partner, and sex workers and their clients. The model is initialized with demographic data from the latest population census, epidemiological data from antenatal care (ANC) surveillance, and behavioral data from the Demographic Health Survey (DHS). The model is first fit to the historical epidemic in the three countries and then used to project the expected number of new HIV infections in the future, with no change in MC levels. We then examined the impact of an expanded program of male circumcision by assuming that the percent of men circumcised would increase to just above 50 percent by 2015 and then remain at this level beyond 2015. Based on recent MC studies, we assumed male circumcision reduces the probability of male infections by 60 percent. We assumed no impact on male-to-female transmission. The difference in the number of new infections between the base projection and the MC program projection represents the reduction in the number of new infections due to scaling up male circumcision.

---

<sup>5</sup> There is no formal agreement on what constitutes a *comprehensive package of MC services*, but for the purposes of this study, it was assumed to contain the following elements: communications, pre- and post behavioral counseling, surgical procedure with post-surgical follow-up, counseling with or without testing, and training.

## Results

Uncomplicated circumcisions usually require four visits: an initial visit for the pre-surgical examination and information and education; a second visit for the surgical procedure; and two follow-up visits at 2–3 and 7 days post-surgery. A fifth visit at 21 days post-surgery is recommended but seldom occurs in uncomplicated cases. Adult MC is done under local anesthesia by all except private providers, who usually use general anesthesia. Waiting time between the first and second visit ranges from 1–8 weeks. Antibiotics are routinely prescribed by some providers but not all. Dressings are generally re-applied at the first post-operative visit (although this was not the case in Swaziland and Zambia). The cost to the patient using non-private providers ranged from US\$4.10 to US\$8.30 in Lesotho and US\$4.84 to US\$41.49 in Swaziland.<sup>6</sup> The unit cost of a comprehensive package of MC services, weighted for the cost of complications, was estimated at \$56.35 (M412) for Lesotho, U\$51.30 (E376) for Swaziland, and U\$46.82 (K200,863) for Zambia. In the three countries, the largest share of this amount was surgical costs (in excess of 80%), followed by communications (approximately a tenth), testing and pre-and post-operative counseling (each less than 5%). Note that these amounts exclude training, community mobilization, and policy analysis and formulation costs, as well as implementation planning and coordination costs.

Table 1 summarizes the results. Based on the cost analysis, the epidemiologic impact and cost-effectiveness of scaling up MC among males (ages 15–49) to 52.5 percent in Lesotho, 57.5 percent in Swaziland, and 58.5 percent in Zambia was calculated. It was estimated that one HIV infection will be averted for every 6 male circumcisions in Lesotho, 4 circumcisions in Swaziland, and 8 circumcisions in Zambia. The cost per infection averted was estimated at US\$292 (M2,136), US\$176 (E1,290) and US\$313 (K1,342,801) for Lesotho, Swaziland, and Zambia, respectively.

Relative to other prevention interventions, MC is potentially a cost-effective intervention. The cost-effectiveness analysis depends on several factors: (1) the period over which the cost-effectiveness analysis is estimated and (2) the pace of scaling up. These findings are largely because MC is a one-time intervention and because there are direct and indirect effects associated with MC. The benefits of male circumcision are therefore multiplicative over time.

**Table I. Summary of Findings**

	Lesotho		Swaziland		Zambia	
Unit Cost	US\$56.35 (M412)		US\$51.30 (E376)		US\$46.82 (K200,863)	
MC Coverage target for 2015	52.5%		57.5%		58.5%	
MCs needed (2008–2020)	Cumulative	Average annual	Cumulative	Average annual	Cumulative	Average annual
	357,143	27,473	150,320	11,297	2,175,896	167,377
Percentage of infections averted relative to baseline	12.9%		10%		13.3%	
Number of infections averted (2008–2020)	Cumulative	Average annual	Cumulative	Average annual	Cumulative	Average annual
	58,931	4,533	36,514	2,809	270,928	20,841
MCs per infection averted	6		4		8	

<sup>6</sup> Information on the cost to the patient using non-private providers was not collected for Zambia.

Cost per infection averted*	US\$292 (M2,136)		US\$176 (E1,290)		US\$313 (K1,342,801)	
Cost of scaling up MC (million)	Cumulative	Average annual	Cumulative	Average annual	Cumulative	Average annual
	US\$17.2 (M125.9)	US\$1.3 (M9.6)	US\$6.6 (E48.3)	US\$0.5 (E3.7)	US\$84.9 (K364,000)	US\$6.5 (K28,000)

How sensitive are the cost-effectiveness results to assumptions about behavioral responses to MC? The impact of changes in condom use on cost per infection averted showed that the results are relatively insensitive to small to moderate reductions. For example, the impact of male circumcision would be less than shown here if those who are circumcised adopt riskier behaviors because they think they are protected by the circumcision. A 25 percent reduction in condom use among circumcised men would reduce the impact by about 7, 17, and 20 percent in Lesotho, Swaziland, and Zambia, respectively. These results underscore the critical importance of (1) locating the surgical provision of MC within a comprehensive set of services that includes behavior change communications and pre- and post-operative counseling; and (2) locating MC services within a broader set of effective prevention interventions.

The scaling up requires approximately an average annual number of 27,473 circumcisions or a daily average of 114 circumcisions in Lesotho through 2020 to achieve and maintain the targeted coverage. In Swaziland, 47 circumcisions are required daily, and in Zambia, 697 circumcisions are required daily through 2020. The human resource implications are important to note. In Lesotho and Swaziland, 10.9 and 6.1 full time physicians are required, respectively. The average annual cost of scaling up MC between 2008 and 2020 in Lesotho is US\$1.3 million (M9.6 million), US\$0.5 million (E3.7 million) in Swaziland, and US\$6.5 million (K28.0 billion) in Zambia. As expected, the surgical procedure accounts for the overwhelming share of the total costs, followed by communications. As mentioned, training, community mobilization, policy analysis and formulation, and implementation planning and coordination costs are not included in these cost estimates.

## Conclusion

The analysis has shown that MC can be a cost-effective intervention when compared with the relative cost effectiveness of other prevention interventions. Several factors influence the potential benefits. The pace of scaling up matters, and the benefits from scaling up are not only multiplicative but also long lasting. In addition, it is important to provide the surgical procedure within a comprehensive set of services that includes behavior change communications and pre- and post-operative counseling and, equally important, to provide MC services within a broader set of effective prevention interventions.

The analysis also showed that the implications to the health system are not trivial. The intention is not to suggest that vast increases in service delivery capacity (in terms of surgical facilities or surgical staff) are necessarily needed in the three countries. Rather, it is recommended that some innovative ways be identified to involve all providers of clinical services. One issue to consider is that public and private provision of MC can be complementary and should not be viewed as mutually exclusive. However, mechanisms should be devised to align practices among private providers with the recommended approaches (routine use of general anesthesia, routine prescription of antibiotics). The positive externalities associated with MC have been established and lay the basis for justifying public subsidization. However, the specific provider payment methods need to be decided on (e.g., fee-for-service, capitation approaches) as well as the mechanisms of reimbursement (vouchers mechanisms, contracting, etc.). The health service, financial, and human resource implications are significant but not insurmountable; taking them into account will be important in ensuring that the benefits of this important public health intervention are realized as soon as possible by the people of Lesotho, Swaziland, and Zambia.



## INTRODUCTION

Clinical trials have now confirmed the efficacy of male circumcision (MC) in reducing female-to-male HIV transmission. Some cost data have been reported (ranging between US\$25 and US\$69), and these cost data also formed the basis of a cost-effectiveness analysis.<sup>7</sup> It is unclear, however, what exactly is included in the costing studies and hence whether these costs are directly comparable. For example, often, indirect costs are not fully reflected; donations (especially clinicians' time) are not costed; variation by provider type and level of health facility is not considered; and the scale of service delivery is unclear. It is anticipated that this cost analysis will provide a more detailed examination of the costs of MC and might inform a sounder basis for an assessment of the cost-effectiveness of MC. This analysis has three components: (1) understanding the social, cultural, and policy context of MC; (2) costing MC; and (3) modeling the impact of MC on the HIV epidemic.

The purpose of this analysis was to assess the cost of providing adult MC in resource-constrained settings, such as Lesotho, Swaziland, and Zambia; and to evaluate the implications of scaling up MC in these countries for the cost-effectiveness of MC as well as for the health system (e.g., resource mobilization and health system capacity).

## METHODOLOGY

### Cost Analysis

A detailed cost analysis of a sub-set of health facilities where circumcision is performed in Lesotho, Swaziland, and Zambia was undertaken. The details of the data collection tool and the costing model are provided in Appendices A and B. The data collection instrument was designed specifically for this study, and the ministries of health can replicate the tool at a later stage to better understand the functioning and key determinants of costs as circumcision is implemented or scaled up.

### Costing Approach

Standard costing methods were applied. The cost data were collected retrospectively through interviews with clinical and administrative staff at health facilities, some nongovernmental facilities, the human resources department of the ministries of health; and from a drug supply structure in the ministry of health in Lesotho. An ingredients approach was used, whereby all the inputs were listed and their contribution to the overall cost was then quantified. The approach allows us to assess whether the costs of the chosen data collection sites can indeed be generalized by making explicit the specific elements of the service that are included in the cost analysis. For example, if a certain cost category is specific to the sites studied, then that item's contribution to the overall unit cost can be removed. This cannot be done if the total program expenditure is merely divided by outputs. This approach is also useful for planning purposes, as it allows planners to add or remove certain elements of the intervention based on the MC guidelines that may be adopted in the respective countries.

*Method of estimation.* The theory behind this estimation is from standard economics theory where total cost ( $C_t$ ) in year,  $t$ , equals price ( $P$ ) multiplied by the quantity ( $Q_t$ ) of circumcisions in any given year,  $t$ .

---

<sup>7</sup> Kahn, J.G., E. Marseille, and B. Auvert. 2006. "Cost-Effectiveness of Male Circumcision for HIV Prevention in a South African Setting." *PLoS Medicine* 3(12).

The aim of this part of the analysis was to estimate the unit cost,  $P_j$ . Unit cost of MC,  $P_j$ , was estimated as:

$$C_j = c_j^{direct} + \sigma * c_j^{indirect}$$

$$c_j^{direct} = \sum_{h,j} c_{h,j}$$

$$c_j^{indirect} = \sum_{l,j} c_{l,j}$$

Where:  $c_{k,j}$  = direct cost per person served at provider,  $j$ , for  $h=1$  (staff), 2 (drugs), 3 (supplies), etc.  
 $c_{l,j}$  = indirect cost per person served at provider,  $j$ , for  $l=1$  (facility equipment), 2 (facility utilities), 3 (facility supervision), etc.  
 $\sigma$  = share of facility services devoted to MC  
Providers ( $j$ ) considered are  $j=1$  (government), 2 (NGO), 3 (missionary)

Direct and indirect costs of MC with complications were also estimated. Unit cost of uncomplicated MCs were weighted by the cost of MC complications and the probability of each complication occurring.

$$C_j^* = pr(\text{uncomplicated MC}) * C_j^{uncomplicated} + pr(\text{complication}_k) * C_j^{complication_k}$$

$$TC_{j,y} = C_j * Q_{j,y}$$

$$TC = \sum_{j,y} \left[ \frac{TC_{j,y}}{(1+r)^a} \right]$$

Where:  $r$  = real discount rate<sup>8</sup>  
 $a$  = number of years into the future

Multiple countries were considered, allowing for standardization, comparison, and validation. The choice of the countries were guided by the two factors that according to Williams and others (2006)<sup>9</sup> were predictors of a large potential benefit from scaling up MC, namely, high HIV prevalence and low to moderate MC prevalence.

**Currency.** The cost data is presented in local currency and in U.S. dollars, using an exchange rate at the time of data collection (January/January/April 2007) (US\$1 = 7.3 Maloti, 7.32 Emalangen, and 4,290.10 Kwacha).

**Scale.** Scale differences may also influence the estimated unit cost. The scale at which MC is implemented at the various sites are of a similar order or magnitude, allowing for comparison of the cost data.

<sup>8</sup> A discount rate of 3 percent was used. This is within the range of 3–5 percent recommended in the World Health Organization’s Guide to Cost-Effectiveness Analysis.

<sup>9</sup> Williams, B.G., J.O. Lloyd-Smith, E. Gouws, Hankins, C., Getz W.M., J. Hargrove, I. De Zoysa, C. Dye, and B. Auvert. 2006. “The Potential Impact of Male Circumcision on HIV in Sub-Saharan Africa.” *PLoS Medicine* 3(7).

*Indirect costs.* The table in Appendix C shows some of the broad categories of costs that were considered. In the costing methodology, goods that have a life of more than one year are treated as capital costs. Financial cost analysis involves estimating the average annual cost of each capital item in terms of simple “straight line” depreciation—the most common form of depreciation.<sup>10</sup> As the table shows, indirect costs also included recurrent costs such as personnel and non-personnel costs.

*Steps in the unit cost estimation.* The steps in the analysis included (1) describe the intervention, (2) identify inputs, (3) quantify inputs, (4) collect cost information, (5) assign monetary values, (6) calculate total cost, (7) quantify outputs, and (8) calculate unit cost.

*Defining the intervention.* Currently, circumcision is not implemented primarily as an HIV prevention intervention; and, therefore, in most instances, not all of the elements of a comprehensive MC package of services were routinely provided. The costing study, however, not only considered the surgical intervention but also the services not currently part of circumcision (information and education, counseling with or without testing, training, etc.); this was done to ensure that the information could be used to inform planning for implementation of a comprehensive package of MC services. Indirect methods were used to get cost data on the non-surgical elements of comprehensive circumcision services. For example, the cost of counseling with or without testing was obtained from voluntary counseling and testing (VCT) programs. In summary, the costing study takes into account the following components shown in Table 1:

**Table 1. Surgical Direct and Indirect Costs and Non-surgical Components**

<b>Surgical Procedure</b>	
<b>Direct Costs</b>	<b>Indirect Costs</b>
Staff	Management and supervision staff
Drugs	Support staff
Consumable supplies	Infrastructure
Non-consumable supplies	Equipment
	Vehicles
	Maintenance
	Utilities
<b>Non-surgical components</b>	
Counseling	
Testing	
Communications	
Training (not included in all the analyses)	

This study focused on the cost of circumcision in adolescent and adult males. Circumcision at birth or in young boys has very different cost implications. For example, circumcision in young boys has to be performed under general anesthesia, and the costs are much higher. On the other hand, neonatal circumcision is a lower cost procedure as the surgical procedure is much simpler and less time consuming, no suturing is required, and minimal post-operative follow-up is needed.

The analysis allows for variation in circumcision practices and patient management. These various clinical practices are made explicit and costed accordingly (e.g., whether there is routine use of antibiotic

<sup>10</sup> Depreciation reflects the loss of value of a capital good over time and involves dividing cost of the item by its useful life or, stated alternatively, by multiplying the cost by the depreciation factor.

prescriptions, whether there is routine use of general anesthesia, etc.)—all of which have important cost implications.

The analysis was undertaken from the perspective of service providers (i.e., not from households or consumers). The implication is that expenses incurred by patients are not included in the analysis (e.g., travel costs, opportunity cost of travel time, opportunity cost of post-operative healing time, etc.). However, this does not mean that the costs facing patients are not taken into account. Data were collected on the out-of-pocket expenses that patients incur for the services from various provider types.

For the surgical intervention, the inputs were quantified from booking to discharge of the patient and all follow-up visits. The inputs included all staff that are involved (clinical, non-clinical staff); time spent by each staff member; salary and non-salary benefits for staff members; and drugs and supplies by type, quantity, and associated costs. In addition to these direct costs, indirect costs were also included, such as facility operating costs (e.g., electricity, maintenance); facility administrative costs (e.g., superintendent, accountant); and facility equipment costs (e.g., autoclaves, refrigerators, vehicles).

*Selection of facilities.* Cost data were collected from public facilities operated by the countries' ministries of health as well as NGO facilities in the three countries. Actual cost data were not collected from private facilities, although private providers were interviewed to obtain information on how they implement MC and the cost to the patient.

### **Epidemiologic Modeling**

Male circumcision has been shown by three randomized control trials to be effective in protecting men from HIV infection. The rate of infection among men circumcised in the trials was about 60 percent lower than the rate among men who were not circumcised. The actual impact in each country will depend on many factors, including the rate of scale up of circumcision services and the dynamics of the epidemic. To estimate the impact that might be expected in Lesotho, Swaziland, and Zambia, we applied a computer simulation model, Spectrum, that can be used to replicate the historical epidemic pattern and demonstrate the effects of increasing the level of male circumcision.

The model simulates the adult population between the ages of 15 and 49 (which accounts for about 85–90% of all adult HIV infections). It is implemented as a module within the Spectrum Suite of Policy Models developed and maintained by the POLICY Project and now the USAID | Health Policy Initiative, Task Order 1. Full details, including the model equations, are provided in the Spectrum user's manual.

The population is divided into male and female populations but is not further stratified by age within the 15–49 age group. New entrants to the model are those people reaching 15 years of age. When they initiate sexual activity, they are placed into one of five risk groups: low-risk heterosexuals (those faithful to a single partner), medium-risk heterosexuals (those with multiple partners), high-risk heterosexuals (sex workers and their clients), men who have sex with men (MSM), and injecting drug users (IDUs). People may leave any of the higher risk groups by adopting low-risk behavior.

Every person entering the model population is assumed to be HIV negative and to remain uninfected while not sexually active. The sexually active and IDU populations are at risk of infection each year. The probability of becoming infected depends on characteristics associated with that individual as well as his or her sexual partners. For sexual transmission, these factors include number of partners, number of acts per partner, condom use, presence of other sexually transmitted infections (STIs), use of antiretroviral therapy (ART), stage of infection, and male circumcision.

Most contact is assumed to be with partners in the same risk group. However, for low-risk groups, contact with other risk groups is a major source of new infection. Therefore, calculations for the low-risk population take into account that some individuals who are faithful to their partners will still be at risk because they have partners who engage in riskier behavior.

A person newly infected with HIV is in the Primary Infection Stage and remains in this category for six months. People in this stage are more infectious than those in other stages. An infected person passes out of the Primary Infection Stage to enter the Asymptomatic Stage, where he/she remains for approximately six years and has a low level of infectiousness. An infected person then moves to the Symptomatic Infection Stage, where he/she remains for about two more years, before dying from an AIDS-related illness. Infectiousness is also elevated for people in the Symptomatic Infection Stage. People are considered to be eligible for ART during this stage. If they receive ART, then their progression to death is reduced.

The probability of becoming infected is modified by male circumcision and other factors. For the analysis described here we assumed that MC reduces the probability of female-to-male transmission of HIV per sexual contact by 60 percent and that there is no effect on the probability of male-to-female transmission. A study is underway in Uganda to test whether male circumcision also reduces male-to-female transmission.

Data for the model are taken from a number of sources. Demographic data are primarily from the United Nations Population Division; HIV prevalence information is from HIV surveillance; and behavioral data come from Demographic and Health Surveys for the three countries. A more detailed explanation of the model is provided in Appendix A. The epidemiology model was linked to the AIDS Impact Module of Spectrum in order to make the HIV projections for the three countries.

## **UNIT COST ESTIMATION**

Table 2–4 summarize the key results of the inputs into the surgical procedure across the facilities of the three countries. Generally, four visits are required. The first visit is for initial examination and booking of the surgical procedure. The nurse and doctor mainly provide information regarding the surgical procedure, healing, potential complications, or adverse events. The waiting time between the first and second visit varies between 1 week and 8 weeks and appears to be constrained by competing surgical needs rather than differences in demand. The second visit involves the doctor, nurse, and nurse’s assistant. The third visit usually happens 2–3 days after the surgical procedure, and the fourth visit is usually 7 days post-surgery. Although a fifth visit at 21 days post-surgery is recommended to the patient, patients with uncomplicated circumcision rarely return for this visit. Staff time spent is shown in each table. The tables also reveal some of the most notable variations among the provider types: (1) the type of anesthesia used, (2) whether antibiotics were routinely prescribed, and (3) whether a surgical dressing was applied after the third visit (at 2–3 days post-surgery).

**Table 2. Summary of Key Results—Lesotho**

<b>Facility</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
Facility type	Hospital	Hospital	Hospital	Hospital
Ownership	MoHSW	MoHSW	CHAL	CHAL
Cost to patient	US\$4.10 (M30 )	US\$4.84 (M35 )	US\$8.30 (M60)	not available
<b>Visit #1 (Initial visit, examination, booking)</b>				
Staff time (minutes)				
Doctor	5	5	10	5
Nurse	10	8	10	5
Counselor	15 <sup>11</sup>	0	0	0
Waiting time	2 weeks	8 weeks	1 week	1 week
<b>Visit #2 (surgical procedure)</b>				
Staff time (minutes)				
Doctor	20	25	39	30
Nurse	20	25	39	30
Nurse Assistant	20	25	20	35
Nurse Anesthetist	0	20	0	0
Anesthesia	Local anesthesia	Local/General anesthesia	Local anesthesia	Local anesthesia
Antibiotics	Cloxacillin	Amoxicillin	Amoxicillin/Cloxacillin	Amoxicillin
Analgesics	Ibuprofen	Ibuprofen	Ibuprofen/Paracetamol	Ibuprofen/Paracetamol
<b>Visit #3 (post-surgery follow-up; +2–3 days)</b>				
Staff time (minutes)				
Doctor	5	8	5	0
Nurse	5	8	0	10
Dressing Re-applied	Yes	Yes	Yes	Yes
<b>Visit #4 (post-surgery follow-up; +7 days)</b>				
Staff time (minutes)				
Nurse	5	5	5	5

<sup>11</sup> Only 30 percent of MC patients.

**Table 3. Summary of Key Results—Swaziland**

<b>Facility</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
Facility type	Hospital	Hospital	Hospital	Health Center
Ownership	MoHSW	MoHSW	NGO/Mission	NGO
Cost to patient	U\$4.84 (E35 )	U\$8.30 (E60 )	U\$4.84 (E35)	US\$41.49 (E300)
<b>Visit #1 (Initial visit, examination, booking)</b>				
Staff time (minutes)				
Doctor	10	10	10	0
Nurse	10	10	10	8
Counselor	0	0	0	28
Waiting time	2 weeks	1 week	8 weeks	1 week
<b>Visit #2 (surgical procedure)</b>				
Staff time (minutes)				
Doctor	18	53	39	50
Nurse	18	53	39	50
Nurse Assistant	18	33	20	33
Nurse Anesthetist	0	0	0	0
Anesthesia	Local anesthesia	Local anesthesia	Local anesthesia	Local anesthesia
Antibiotics	None	None	None	None
Analgesics	Paracetamol	Paracetamol	Paracetamol	Paracetamol
<b>Visit #3 (post-surgery follow-up; +2–3 days)</b>				
Staff time (minutes)				
Doctor	15	15	15	0
Nurse	0	0	0	15
Dressing Re-applied	No	Yes	Yes	No
<b>Visit #4 (post-surgery follow-up; +7 days)</b>				
Staff time (minutes)				
Nurse	5	5	5	5

**Table 4. Summary of Key Results—Zambia**


<b>Facility</b>	<b>1</b>	<b>2</b>	<b>3</b>
Facility type	Hospital	Hospital	Clinic
Ownership	Public	Public	NGO/Private
<b>Visit #1 (initial visit, examination, booking)</b>			
Staff inputs (minutes)			
Nurse	30	37	37
<b>Visit #2 (surgical procedure)</b>			
Staff inputs (minutes)			
Nurse	30	28	28
Clinical Officer	14	5	5
Post graduate doctor	14	0	23
Urologist	3	0	0
Anesthetist	0	0	5
Type of Anesthesia	Local anesthesia	Local anesthesia	Local anesthesia
Antibiotics	None	None	None
Analgesics	Paracetamol	Paracetamol	Paracetamol
<b>Visit #3 (post-surgery follow-up; +2 days)</b>			
Staff inputs (minutes)			
Nurse	15	37	37
Clinical Officer	10	5	0
Consultant Urologist	11	0	0
Anesthetist	0	0	5
Dressing Re-applied	No	No	No
<b>Visit #4 (follow-up; +7 days)</b>			
Staff inputs (minutes)			
Nurse	5	5	5
Clinical Officer	5	5	0



## Direct and Indirect Costs


Figure 1 shows the variation in direct costs and indirect costs for the facilities visited. There was general convergence in the level of the direct costs, but the indirect costs showed substantial variation across facilities. The breakdown of the direct and indirect costs associated with an uncomplicated MC for the three countries is shown in Figure 2. In all three countries, the largest contributors to the overall costs were staff and consumable supplies. In Lesotho, maintenance and utilities were also major contributors to the cost.

**Figure 1. Variation in Direct and Indirect Costs Across Facilities**




**Figure 2. Breakdown of Cost of Circumcision**


**Lesotho**



**Swaziland**



**Zambia**



## Complications and Related Costs

Post-operative complications are relatively uncommon occurrences, occurring in just over a tenth of circumcisions in the facilities studied. Mild to moderate pain is generally not viewed as a complication, and analgesics are routinely prescribed as part of the procedure. The most common post-operative complications<sup>12</sup> are hemorrhage and/or hematoma and sepsis. These are complications of moderate severity, and no cases of the other adverse events with long-term consequences described in the literature<sup>13</sup>—such as excessive skin removal, problems with urination, or problems with appearance—were reported. Hemorrhage and/or hematoma are the most resource intensive complications, as they require anesthesia, re-opening of the surgical wound, and ligation of each bleeding vessel individually. Hemorrhage is usually detected shortly after surgery and sometimes even before the patient is discharged. Hematoma results from internal bleeding and commonly involves swelling; the patient usually returns within the first day after the surgery. The other complication, sepsis, is usually detected at the first post-operative visit (2–3 days after surgery).

The two most common complications were costed (hemorrhage and sepsis). These are reflected in an adjusted unit cost weighted for the additional cost of complications and the probability of each complication occurring. Figure 3 shows a breakdown of unit costs of MC without complication, unit costs if a complication arises, and the adjusted (weighted) unit cost. The unit cost of MC complicated by hemorrhage is between 22.3 and 57.1 percent higher than an uncomplicated MC. For MC complicated by sepsis, the cost is between 7.1 and 18.5 percent higher. Because of the relative infrequent occurrence of the complications, the unit cost weighted by the frequency of complications is only between 1.9 and 3.3 percent higher than a circumcision without any complication. The most important sources of the difference are the cost of consumable supplies and staff costs.

The results presented here show the inputs and associated costs of circumcision as it is currently implemented in the three countries. There are important omissions that are routinely included as part of a complete MC package of services (e.g., communications, testing, pre- and post-circumcision information and education, and behavioral counseling and training). As mentioned in the description of the methodology, the costs for the additional interventions (pre- and post-circumcision behavioral counseling and HIV testing) were taken from existing programs that offer these services but might be unrelated to MC. The assumptions that underpin the estimation of these costs are shown in Appendix D.


In the comprehensive package of MC services pre- and post-circumcision behavioral counseling, HIV testing, and communications collectively add between 24.0 and 30.7 percent to the cost of the surgical procedure. Communications account for the largest share of this additional cost. Training costs were also estimated according to the assumptions listed in Appendix D, but these costs have not been included in Figure 2 and Figure 3. This is largely because training needs are dependent on the human resource requirements and do not necessarily have to be repeated annually, and additional consultation is needed to agree on training needs. Other excluded costs are policy analysis and formulation and community mobilization costs.

---

<sup>12</sup> Note that this is not a comprehensive survey of the complications associated with MC. These figures are merely used to apply probability weights to the cost data collected. A more detailed assessment of MC complications has been conducted for Swaziland, and the rate of MC complications was lower. The differences may be due to differences in definitions and the period over which the complications were studied.

<sup>13</sup> UNAIDS, WHO, London School of Hygiene and Tropical Medicine. Draft 2007. Male circumcision: Global Trends and Determinants of Prevalence, Safety, and Acceptability.

**Figure 3. Comparison of the Breakdown of Adjusted and Unadjusted Unit Costs of Surgical Procedure**



## SCALING UP CIRCUMCISION

The future uptake and rate of scale-up of MC is not known. However, to inform planning processes, the results of scenario modeling are presented.

The scenarios assume that 5, 15, and 16.9 percent of males are effectively circumcised (i.e., circumcised with sufficient removal of foreskin to afford the protective effect found in the clinical trials) in Lesotho, Swaziland, and Zambia, respectively. The projection furthermore aims to achieve coverage of 50, 57.5, and 58.5 percent of adult males in the three countries, respectively, by 2015. A linear scale-up is assumed. The proportion of men who are HIV positive is not factored into this scenario because the MC protocols drafted to date (i.e., by the WHO) do not specify the requirement of an HIV-negative status for MC eligibility. Because MC is a one-time procedure, the number of MCs that need to be performed in a particular year is the difference between coverage estimates for that year and the previous. The number of circumcised males that exit the age group and uncircumcised males that enter this age group are also taken into account.

### Impact and Cost-effectiveness of Scaling up MC

According to the baseline epidemiologic model, the cumulative number of new infections over the period 2008–2020, given the current level of behavior, will be 455,589 or on average 35,045 annually in Lesotho; 373,836 or on average 28,757 annually in Swaziland; and 1.95 million or on average 150,028 annually in Zambia. Table 5 presents the results of the analysis if MC were to be scaled up. It is projected that cumulatively, over the period 2008–2020, 58,931 new infections will be averted in Lesotho, 36,514 in Swaziland, and 270,928 in Zambia. This implies that between 8.9 and 13.3 percent of new infections will be averted relative to the baseline projection in the three countries. The implication is that, on average, over the year period 2008–2020, 1 new HIV infection will be averted for every 6 circumcisions in Lesotho, 4 in Swaziland, and 8 in Zambia. Using the cost per MC from the cost analysis, the discounted cost per infection averted is estimated at \$292 (M2,136) in Lesotho, US\$176 (E1,290) in Swaziland, and US\$313 (K1,342,801) in Zambia.

**Table 5. Impact and Cost-Effectiveness of Scaling Up MC (2008–2020)**

	Lesotho		Swaziland		Zambia	
	Cumulative	Average annual	Cumulative	Average annual	Cumulative	Average annual
Number of infections averted	58,931	4,533	36,514	2,809	270,928	20,841
Percentage of infections averted relative to baseline	12.9%		10%		13.3%	
MCs per infection averted		6		4		8
Cost per infection averted*		US\$292 (M2,136)		US\$176 (E1,290)		US\$313 (K1,342,801)

\*discounted

How does this compare with the cost-effectiveness of other prevention interventions? Table 6 shows the results from literature on the cost-effectiveness of prevention interventions and suggests that at a cost per infection averted of between US\$176 and US\$313, MC is a highly cost-effective intervention. This estimate is also consistent with other studies on the cost-effectiveness of MC.

**Table 6. Cost per HIV Infection Averted for Selected HIV Interventions<sup>14</sup>**


<b>Intervention</b>	<b>Cost per Infection Averted</b>
Condom distribution	US\$10–US\$2,188
VCT	US\$393–US\$482
PMTCT	US\$20–US\$2,198
STI treatment	US\$271–US\$514
School-based education	US\$7,288–US\$13,326
Male circumcision	US\$181

## **Pace of Scale-up**

The preceding analysis assumed a linear scale-up to reach the target coverage by 2015. What if the pace of scaling up moves slower or faster than a linear scale-up, as illustrated in Figure 4? Table 7 shows the impact of various patterns of scale-up relative to the linear scale-up. If MC scale-up occurred at a slower pace, the average annual number of MCs over the period 2008–2020 would be similar, but the number of infections averted would be much less (between -19.7% and -14.5%) relative to the linear scale-up. If, however, scale-up occurred at a faster pace, as illustrated in Figure 4, the average annual number of MCs needed would again be similar, but the number of infections averted would be higher (between 13.7% and 16.1%) relative to the linear scale-up. As expected, the number of MCs needed to avert one HIV infection is lower for the faster scale-up and the cost per infection averted is lower by between 20 and 33 percentage points relative to the slow scale-up scenario. The reason for these findings relate to the indirect effects associated with MC and the resultant multiplicative impact of MC illustrated in Figure 5.

<sup>14</sup> Creese A, K. Floyd, A. Alban, and L. Guinness. 2002. “Cost-effectiveness of HIV/AIDS Interventions in Africa: A Systematic Review of the Evidence.” *Lancet* 2002 (359):1635–43. Kahn, J.G., E. Marseille, and B. Auvert. 2006. “Cost-Effectiveness of Male Circumcision for HIV Prevention in a South African Setting.” *PLoS Medicine* 3(12).

**Figure 4. Alternate Patterns of Scaling Up MC**




**Table 7. Impact of Various Pace of Scaling Up MC (2008–2020)**

<b>Lesotho</b>	<b>Linear scale-up</b>	<b>Slower scale-up</b>	<b>Faster scale-up</b>
	2008–2020	% difference with linear scale-up	
Average annual number of MCs	27,473	-1.6%	+1.5%
Average annual number of infections averted	4,533	-16.4%	+15.9%
MC per infection averted	6	+13.6%	-9.6%
Cost per infection averted*	US\$292 (M2,136)		
<b>Swaziland</b>	<b>Linear scale-up</b>	<b>Slower scale-up</b>	<b>Faster scale-up</b>
	2008–2020	% difference with linear scale-up	
Average annual number of MCs	11,297	-1.2%	+1.3%
Average annual number of infections averted	2,809	-14.5%	+13.7%
MC per infection averted	4	+12.1%	-8.2%
Cost per infection averted*	US\$176 (E1,290)		
<b>Zambia</b>	<b>Linear scale-up</b>	<b>Slower scale-up</b>	<b>Faster scale-up</b>
	2008–2020	% difference with linear scale-up	
Average annual number of MCs	167,377	-1.1%	+0.09%
Average annual number of infections averted	20,841	-19.7%	+16.1%
MC per infection averted	8	+19.3%	-11.0%
Cost per infection averted*	US\$313 (K1,342,801)		

\*discounted

**Figure 5. Indirect Effects Associated with MC**



### **Multiplicative Impact of MC**

Figure 5 presented by Hallet and others (2007)<sup>15</sup> show the direct and indirect effects associated with MC. The direct effect is the benefit that accrues to a circumcised male and has been the subject of the MC clinical trials. The various indirect effects are determined by the epidemiological dynamics. First-, second-, and third-line indirect effects are illustrated in Figure 5. Because of the protective effect of circumcision, fewer circumcised men will become infected over time. The secondary result is that female sexual partners of these circumcised men are less likely to become infected. This is the first-hand indirect effect. The second-hand indirect effect comes about because these women are less likely to infect other males, whether they are circumcised or not. Following on the second hand indirect effect, uninfected women are also less likely to be infected, resulting in the third hand indirect effect.

Note that not all the benefits implied in the cost-effectiveness results will be realized within the first few years of implementation. Figure 6 shows the annual estimates of the cost per infection averted of MC, starting at a high of between US\$1,661 and US\$3,036 per infection averted in 2008 and decreasing to between US\$35 and US\$77 per infection averted in 2020. This result is derived from the fact that the percent of new infections averted thanks to scaling up MC is initially only between 1.5 and 1.7 percent in 2008 and increases to between 13.0 and 21.8 percent in 2020.


---

<sup>15</sup> Presentation at HIV Implementers' Conference, Kigali, June 2007. "Understanding the Impact of Male Circumcision as an Intervention" Authors: Smith J., T. Hallett, S Gregson, B Lopman, K Desai, M. Boily, G. Garnett, Department of Infectious Disease Epidemiology, Imperial College London. Presenter: Tim Hallett.




**Figure 6. Estimates of Cost per Infection Averted by Year**


**Lesotho**



**Swaziland**



**Zambia**



## Implications to the Health System

Table 8 shows some of the health service implications of scaling up MC among adult males (15–49 years old). It is estimated that scaling up requires an average annual number of 27,473 circumcisions in Lesotho, 11,297 in Swaziland, and 167,377 in Zambia. The implied daily average is 114 circumcisions in Lesotho, 47 in Swaziland, and 697 in Zambia through 2020 to achieve and maintain targeted coverage.

**Table 8. Health Service Implications**

	<b>2008–2020 Cumulative</b>	<b>2008–2020 Average Annual</b>
<b>Lesotho</b>		
# of new circumcisions needed	357,143	27,473
# of circumcisions per month		2,289
# of circumcisions per week		572
# of circumcisions per day		114
<b>Swaziland</b>		
# of new circumcisions needed	150,320	11,297
# of circumcisions per month		941
# of circumcisions per week		235
# of circumcisions per day		47
<b>Zambia</b>		
# of new circumcisions needed	2,175,896	167,377
# of circumcisions per month		13,948
# of circumcisions per week		3,487
# of circumcisions per day		697

Based on the data collected in the costing analysis, the facilities currently implementing MC perform between 20 and 30 circumcisions per month. Currently, MC is performed exclusively at hospitals, and this implies that if the current approach is followed for scaling up MC, a substantial increase in the number of facilities offering MC is needed at the present level of service delivery. Table 9 shows the estimated number of full-time equivalents (FTEs) per staffing category (needed over the period 2008–2020) that have to be dedicated full-time to male circumcision.


**Table 9. Human Resource Implications**

	<b>2008–2020 Average Annual Staff FTEs Needed</b>
<b>Lesotho</b>	
Counselor	11.9
Surgical nurse	9.1
Nurse assistant	7.1
Physician	10.9
Lab technician	2.4

<b>Swaziland</b>	
Counselor	2.8
Surgical nurse	4.5
Nurse assistant	2.5
Physician	6.1
Laboratory technician	1.0
<b>Zambia</b>	
Surgical nurse	157
Clinical officer	22
Post-graduate doctor	27
Consultant urologist	24

Figure 7 shows the resource implications of MC scale-up. The average annual cost for scaling up MC in Lesotho, Swaziland, and Zambia is US\$1.3 million (M9.6 million), US\$0.5 million (E3.7 million), and US\$6.5 million (K28.0 million), respectively (see Figure 7). As mentioned before, training, community mobilization, and policy analysis and formulation costs are not included in this cost estimate.

**Figure 7. Cost Implications of Scaling Up MC**



\*Size of circles denote cost per infection averted

## Behavioral Responses to MC

How sensitive are the cost-effectiveness results to assumptions about behavioral responses to MC? The impact of changes in condom use on cost per infection averted showed that the results are relatively insensitive to small to moderate reductions. For example, the impact of male circumcision would be less than shown here if circumcised men adopt riskier behaviors because they think they are protected by the circumcision. A 25 percent reduction in condom use among circumcised men would reduce the impact by about 7, 17, and 20 percent in Lesotho, Swaziland, and Zambia, respectively. These results underscore the critical importance of (1) locating the surgical provision of MC within a comprehensive set of services that includes behavior change communications and pre- and post-operative counseling, and (2) locating MC services within a broader set of effective prevention interventions.

## **Sustainability**

MC is a one-time intervention, and, as discussed, the benefits are not only multiplicative but also long lasting. One key driver of the annual costs estimated is the circumcision of 15-year-olds as they enter the 15–49-year-old age group. While neonatal MC was not the focus of this analysis, it is widely acknowledged that neonatal MC is a lower cost procedure for various reasons: the surgical procedure is much simpler and less time consuming, no suturing is required, minimal post-operative follow-up is needed; and finally, the potential for risk compensation is reduced. Simultaneous scaling up of adult and neonatal MC offers an opportunity for some long-term cost savings and sustainability of the gains in preventing HIV; reduced numbers of adult MCs would be needed beyond 2023. However, the scaling up of MC has two important challenges: (1) the cultural context of MC, particularly where MC is part of the right of passage to adulthood, and (2) the small number of births delivered in health service settings where neonatal MC can be performed.

## **Limitations**

Some limitations of the study are worth noting.

- The cost analysis of the non-surgical elements of comprehensive MC services (e.g., counseling, communications, testing, etc.) was not as extensive as the analysis of the surgical procedure. However, the analysis of the non-surgical elements may be refined as a more detailed approach to implementation becomes clear.
- The analysis only considered the impact of MC on reducing HIV transmission and the number of HIV infections averted. It is, however, increasingly being realized that the scaling up of MC offers an opportunity to improve the delivery of male reproductive health services. A similar analysis of the costs and benefits of MC within the context of a comprehensive package of male reproductive health services might be useful.
- Male circumcision is culturally practiced in Lesotho and Zambia, but the analysis did not take into account the cost of MC provision in non-facility settings. Rather, the study focused on male circumcision services provided in government and nongovernmental health facilities in order to provide key decisionmakers and policymakers with the financial resource requirements for scaling up MC within the health system.

## **CONCLUSION**

Male circumcision can be a cost-effective intervention when compared with the relative cost-effectiveness of other prevention interventions. Several factors influence the potential benefits. The analysis showed that the pace of scaling up matters and that the benefits from scaling up are not only multiplicative but also long lasting. The analysis also illustrated the importance of locating the surgical provision of MC within a comprehensive set of services that includes behavior change communications and pre- and post-operative counseling and, equally important, locating MC services within a broader set of effective prevention interventions.

The human and financial resource implications of scaling up MC are significant. The intention is not to suggest that vast increases in service delivery capacity (in terms of surgical facilities or surgical staff) are necessarily needed in Lesotho. Rather, it is recommended that some innovative ways be identified to involve all providers of clinical services. One issue to consider is that public and private provision of MC

can be complementary and should not be viewed as mutually exclusive. However, mechanisms should be devised to align practices among private providers with the recommended approaches (routine use of general anesthesia, routine prescription of antibiotics). The positive externalities associated with MC have been established and lay the basis for justifying public subsidization. However, the specific provider payment methods need to be decided on (e.g., fee-for-service, capitation approaches), as well as the mechanisms of reimbursement (voucher mechanisms, contracting, etc.). The health service, financial, and human resource implications are significant but not insurmountable; taking them into account will be important in ensuring that the benefits of this public health intervention are realized as soon as possible by the people of Lesotho, Swaziland, and Zambia.

## APPENDIX A. EPIDEMIOLOGY MODELING

The analysis used the Spectrum suite of policy models. The various components of Spectrum make it uniquely suited to address the policy questions raised in this study. The *HIV transmission model* is a deterministic model that reveals the key processes or risk factors that give rise to the sexual and intravenous transmission of HIV.<sup>16</sup> The model divides the population into multiple risk groups or compartments, reflecting the heterogeneity in sexual activity (frequency of sex, number of sexual partners, anal sex between males, etc.) and allows for movement (mixing) between the risk compartments. The model is fundamentally based on the Weinstein equation,<sup>17</sup> takes into account heterosexual and homosexual transmission, and is applied to the various risk groups. Example factors that the model takes into account are

- Changes in risk behaviors over time to reflect the impact of prevention interventions;
- Various levels of probability of infection based on the presence of ulcerative and non-ulcerative STIs; and
- Various probabilities of HIV transmission based on the stage of infection (primary infection, asymptomatic, symptomatic) of the HIV-positive individual.

The key variables used for the transmission model are

$$P(\text{Infection})_{r,s,c} = 1 - \left\{ p_r \sum_{s,c} w_{s,c} \left[ 1 - q_{g,s,c} (1 - f_r e) g \right]^{n_r} + (1 - p_{rp}) \right\}^{m_r}$$

r = risk group

s = STI status

c = circumcision status

p = HIV prevalence in partner

q = probability of HIV transmission by partner combination, STI status, and male circumcision status

f = condom use

e = condom efficiency

g = multiplier based on the distribution of infected population by stage of infection

n = number of contacts per sexual partner

m = number of sexual partners

The transmission probabilities used in the model represent the best estimate based on the literature. These values are periodically reviewed and updated to ensure that they reflect the most recent literature. These values are stated explicitly and are also used in sensitivity analyses.

Although not all the factors considered by the model are directly affected by male circumcision (e.g., transmission between male sexual partners), it is essential to comprehensively and accurately model the HIV epidemic in each country before looking at the impact of circumcision.

International agencies involved with setting guidelines and norms (WHO, UNAIDS, etc.) have stressed the need to view male circumcision as part of a comprehensive prevention strategy (i.e., that the biologically based interventions should be complementary to the behaviorally based interventions). The methodology proposed here is uniquely suited to analyze the impact of male circumcision alongside other HIV interventions.

---

<sup>16</sup> While not discussed here, other forms of transmission are reflected in the model. The model treats intravenous transmission similar to the way sexual transmission model is treated here. The model also takes into account other forms of HIV transmission (mother-to-child transmission and infected blood products).

<sup>17</sup> Weinstein, M.C., J.D. Graham, J.E. Siegel, and H.V. Fineberg. 1989. "Cost-Effectiveness Analysis of AIDS Prevention Programs: Concepts, Complications, and Illustrations." In Turner, C.F., H.G. Miller, L.E. Moses (eds.). 1989. *Confronting AIDS: Sexual Behavior and Intravenous Drug Use* 1989: 471–499. Washington, DC: National Academy Press.

## APPENDIX B. DATA COLLECTION INSTRUMENT

### FACILITIES DATA COLLECTION FORM

Name of Facility: \_\_\_\_\_

District: \_\_\_\_\_

Local currency

Exchange rate

Type of Facility:     Health Post     Health Centre     Hospital

Facility Ownership:     Private NGO     Private Commercial     Public

Average Number of Clients  
at this Facility per Months:

Average Number of Male Circumcision  
Contacts at this Facility per Month:

Name and Location of Nearest Health  
Centre (for Health Posts only): \_\_\_\_\_

Distance from Facility:

Name and Location of Nearest Hospital: \_\_\_\_\_

Distance from Facility:

## I. STAFFING, EQUIPMENT, AND MAINTENANCE

### A. CLINICAL STAFF

#### Number of Full-Time Employees

	Number of full-time employees	Annual salary (including benefits)	% of time spent on male circumcision	Notes
Auxiliary/Attendant				
Nurse/Midwife				
General Physician				
Surgeon				
Paediatrician				
Anaesthetist				
Lab Technician				
Counselor				

### B. SUPPORT STAFF

#### Number of Full-Time Employees (*entire facility, not only for male circumcision*)

	Number of full-time employees	Annual salary (including benefits)	Notes
Guard			
Housekeeping			
Reception			
Records			
Supply Clerk			
Maintenance			
Mgmt Officer			
Driver			
Food Preparer			

### C. EQUIPMENT

#### Equipment for *Entire Facility*

	Number of items at facility	Amortization period	Estimated construction/purchasing price	Notes
Operating theatre				
Anaesthesia equipment				
Laboratory				
Sterilizer				
Refrigerators				



Portable refrigerators/cooling boxes				
Furniture/beds				
Overall facility/structure				

**D. EMERGENCY TRANSPORTATION VEHICLES**

**Equipment for *Entire Facility***

	Number of items at facility	Amortization period	Estimated purchasing price	% of time used for transportation of male circumcision complications (estimated)	Notes
Bicycle					
Motorcycle					
Jeep					
Ambulance					

**E. MAINTENANCE AND UTILITY COSTS**

**Equipment for *Entire Facility***

	Annual cost	Notes
Building maintenance		
Utilities Cost		
Emergency Vehicle Maintenance		

## II. CIRCUMCISION PROCEDURE, INCLUDING COUNSELING & TESTING

### A. GENERAL

Provided here       Not provided here, referred to: \_\_\_\_\_

1. Average number of MC clients each month:


2. How many visits does an MC client have on average to obtain both counselling and testing and the procedure?


3. Is informed consent obtained (risks/benefits of procedure, other ways to reduce risk of HIV infection)?


### B. DRUGS, SUPPLIES, AND CONSUMABLES

4. What supplies are used? Are there any other drugs/supplies they receive? (*Write in*)

Drug or Supply	% of MC clients receiving this drug/supply	Number	Unit cost	Notes
Sutures				
Needles				
Bandages				
Analgesic				
Gloves				
Antiseptic				
Gentian violet/sterile marker pen				

5. What percentage undergo a physical exam?

--

### C. LAB TESTS

*Skip if there are no lab facilities:*

6. What percentage of MC clients receive the following lab tests during their circumcision visits?

Lab Tests	% of MC clients receiving this test	Number	Unit Cost	Notes
RPR Syphilis				
HIV				

## II. CIRCUMCISION PROCEDURE, INCLUDING COUNSELING & TESTING (Page 2)

### D. STOCK-OUTS

7. Have you experienced stock-outs, or difficulties with obtaining supplies, with any of the supplies listed above? If yes, how often?

--

### E. CLINICAL STAFF TIME

8. Who provides counselling prior to the circumcision procedure at the facility? How long does the average visit last? Who provides circumcision at the facility? How long does the average visit last?

Pre-surgical visit	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Surgical Visit	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

### F. COMMENTS

--

### III. POST-CIRCUMCISION CARE—NORMAL

#### A. GENERAL

Provided here       Not provided here, referred to: \_\_\_\_\_

1. Average number of post-MC clients each month:       2. How many post-care visits does an MC client have on average?

3. What is included as part of post-circumcision care?

#### B. DRUGS, SUPPLIES, AND CONSUMABLES

4. What supplies are used? Are there any other drugs they receive? (*Write in*)

Drug or Supply	% of MC clients receiving this drug/supply	Number	Unit cost	Notes

5. What percentage undergo a physical exam?       6. Are gloves used?  Yes  No

#### C. LAB TESTS

*Skip if there are no lab facilities:*

7. What percentage of MC clients receives the following lab tests during their follow-up visits?

Lab Tests	% of MC clients receiving this test	Number	Unit cost	Notes

### III. POST-CIRCUMCISION CARE—NORMAL (Page 2)

#### D. CLINICAL STAFF TIME

8. Who provides post-circumcision care at the facility? How long does the average visit last?

Post-surgical visit #1	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Post-surgical visit #2	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Post-surgical visit #3	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

#### E. COMMENTS

## IV. POST-CIRCUMCISION CARE—COMPLICATIONS

### A. GENERAL

Provided here       Not provided here, referred to: \_\_\_\_\_

1. Average number of post-MC clients requiring management of complications each month:       2. How many post-care visits does an MC client with complications have on average?

3. In your opinion, what percentage of patients do you think return to the facility for treatment of complications?

4. What percentage of patients do you treat for MC-related complications who did not receive the procedure at this facility?

### B. DRUGS, SUPPLIES, AND CLINICAL STAFF TIME

4. What percentage of MC clients experience infection?

5. What drugs and supplies are used to treat this complication? (*Write in*)

Drug or Supply	% of MC clients receiving this drug/supply	Number	Unit cost	Notes

6. Who provides care for this complication at the facility? How long does the average visit last?

Visit #1 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Visit #2 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Visit #3 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

7. What percentage of MC clients experience **excessive bleeding**?

8. What drugs and supplies are used to treat this complication? (*Write in*)

Drug or Supply	% of MC clients receiving this drug/supply	Number	Unit cost	Notes

9. Who provides care for this complication at the facility? How long does the average visit last?

Visit #1 for the complication	No. of minutes spent with client	Visit #2 for the complication	No. of minutes spent with client	Visit #3 for the complication	No. of minutes spent with client
Counselor		Counselor		Counselor	
Nurse/Midwife		Nurse/Midwife		Nurse/Midwife	
General Physician		General Physician		General Physician	
Surgeon		Surgeon		Surgeon	
Auxiliary/Attendant		Auxiliary/Attendant		Auxiliary/Attendant	
Anaesthetist		Anaesthetist		Anaesthetist	
Lab Technician		Lab Technician		Lab Technician	
Other		Other		Other	

10. What percentage of MC clients experience **excessive pain**?

11. What drugs and supplies are used to treat this complication? (*Write in*)

Drug or Supply	% of MC clients receiving this drug/supply	Number	Unit cost	Notes

12. What **other complication** do MC clients experience? (*Write in*)

13. What percentage of MC clients experiences this complication?

14. What drugs and supplies are used to treat this complication? (*Write in*)

Drug or Supply	% of MC clients receiving this drug/supply	Number	Unit cost	Notes

15. Who provides care for this complication at the facility? How long does the average visit last?

Visit #1 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Visit #2 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Visit #3 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

16. What **other complication** do MC clients experience? (*Write in*)

17. What percentage of MC clients experiences this complication?

18. What drugs and supplies are used to treat this complication? (*Write in*)

Drug or Supply	% of MC clients receiving this drug/supply	Number	Unit cost	Notes



19. Who provides care for this complication at the facility? How long does the average visit last?

Visit #1 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Visit #2 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

Visit #3 for the complication	No. of minutes spent with client
Counselor	
Nurse/Midwife	
General Physician	
Surgeon	
Auxiliary/Attendant	
Anaesthetist	
Lab Technician	
Other	

**C. LAB TESTS**

*Skip if there are no lab facilities:*

20. Do MC clients with complications receive any lab tests?

Lab Tests	% of MC clients receiving this test	Number	Unit cost	Notes

**D. COMMENTS**

## APPENDIX C. EXAMPLES OF INDIRECT COSTS

<b>Capital costs</b>	
Equipment	Other
Autoclave	Beds
Dental equipment	Office furniture
Examination & surgery equipment	Generator and incinerator
Physiotherapy equipment	Pharmacy equipment
<b>Recurrent costs</b>	
Personnel	Non-personnel
Management	Building maintenance cost
District hospital administrator	Utilities cost
Health inspector	Emergency vehicle maintenance
Asst Health inspector	
Support staff	
Senior pharmacy technician	
Pharmacy technician	
Lat technologist	
Assistant HR officer	
Technical officer	
Senior health assistant	
Health assistant	
Radiographic assistant	
Lab assistant	
Driver	
Accountant	

## APPENDIX D. ASSUMPTIONS USED FOR UNIT COST ESTIMATION OF COMPREHENSIVE PACKAGE OF SERVICES

<b>Communications</b>				
<b>Counseling</b>				
	<i>Pre-Circumcision</i>	<i>Circumcision</i>	<i>Post-Circumcision</i>	<i>Circumcision Follow-up</i>
Time spent by counselor (minutes)	20	10	10	10
<b>Testing</b>				
	<i>Data Inputs</i>			
Commodity costs				
ELISA test	\$2.34			
Rapid test	\$2.46			
Share of tests that are Rapid Tests	80%			
subtotal	\$2.43			
Laboratory technician costs				
Staff time per test (minutes)	10			
Cost of staff time per test	\$1.26			
Share of MC clients tested	50%			
Contribution to Unit Cost (unweighted)	\$3.69			
Contribution to Unit Cost	\$1.85			
Lab technician staff time per test (minutes)	10			
<b>Training</b>				
<i>Training of surgical staff</i>				
Number of training days (doctors)	1			
Number of training days (nurses)	2			
Number of doctors trained	20			
Number of nurses trained	40			
Number of staff days (doctors)	20			
Number of staff days (nurses)	80			
Cost of workshop				
Cost per person per training day	\$68			
Number of training days	100			
Subtotal	\$6,831			
Cost of staff time				
Hours of doctors time	160			
Hours of nurses time	640			
Cost of doctors staff time	\$1,476			
Cost of nurses staff time	\$4,841			

Total	\$13,147
<i>Training of counselors</i>	
Number of training days (counselors)	3
Number of counselors trained	50
Number of staff days (counselors)	150
Cost of workshop	
Cost per person per training day	\$68.31
Number of training days	150
Subtotal	\$10,246
Cost of staff time	
Hours of counselors time	1,200
Cost of counselors staff time	\$2,447
Total	\$12,693
<b>Total Annual Training Costs</b>	<b>\$25,840</b>
<b>Total Annual Training Costs (excl staff costs)</b>	<b>\$17,076</b>

## APPENDIX E. EQUIPMENT REQUIREMENTS

<b>Circumcision Surgical Tray</b>	<b>Quantity</b>	<b>Unit cost (\$)</b>
Gallipot	1	8.05
Sponge holding forceps 18cm	1	69.75
Bistouri scalpel blade holder #4	1	14.80
Straight mosquito artery forceps small	2	13.40
Curved mosquito artery forceps small	2	13.40
ForcepsN/H mayo hager 14–16cm	1	19.50
XAdson fine non-toothed dissecting forceps	1	16.00
Surgical scissors BL ST	1	15.40
Mayo scissors	1	24.60
Dissecting tray and lid	0	123.75
<b>Emergency Tray</b>	<b>Quantity</b>	<b>Unit cost (\$)</b>
Solu Cortef 100mg/2ml	1	27.34
Adrenaline 1mg/2ml	1	15.50
Atropine 1mg/ml	1	18.90
Diazepam 10mg/2ml	1	23.00
IV canula (Jelo radiopaque) 18 guage	1	12.20
Solution administration set	1	25.00
Sodium Chloride 0.9% 1L	1	9.60
Velcro tourniquets	1	10.65

Source: Dr. Adam Groeneveld, Ministry of Health and Social Welfare, Mbabane, Swaziland.

Health Policy Initiative, Task Order I  
Constella Futures  
One Thomas Circle, NW, Suite 200  
Washington, DC 20005 USA  
Tel: (202) 775-9680  
Fax: (202) 775-9694  
Email: [policyinfo@healthpolicyinitiative.com](mailto:policyinfo@healthpolicyinitiative.com)  
<http://ghiqc.usaid.gov>  
<http://www.healthpolicyinitiative.com>