



Fact Sheet on Youth Reproductive Health Policy

GENDER-BASED VIOLENCE

Various Forms of GBV include battering, intimate partner violence (including marital rape, sexual violence, and dowry-related violence), female infanticide, femicide, sexual abuse of female children in the household, honor crimes, early marriage, forced marriage, female genital cutting and other traditional practices harmful to women, sexual harassment in the workplace and educational institutions, commercial sexual exploitation, trafficking of girls and women, and violence perpetrated against domestic workers (USAID, 2009).

“Gender-based violence (GBV) is violence involving men and women, in which the female is usually the victim; and which is derived from unequal power relationships between men and women. Violence is directed specifically against a woman because she is a woman, or affects women disproportionately. It includes, but is not limited to, physical, sexual, and psychological harm. It includes that violence which is perpetuated or condoned by the state” (UNFPA, 1998).

GBV encompasses a range of physical, sexual, and psychological violence that disproportionately affects women and girls (IGWG/USAID, 2008). Violence against women is a major problem that negatively affects the reproductive health of women and girls. “Each year as many as 40 million children under 15 years old experience some form of violence. Many of them are girls, and many live in Africa” (African Child Policy Forum, 2006).

Although the global community has focused greater attention on GBV in recent years, levels of violence against women remain unchanged (UN General

Assembly, 2006a). Eliminating GBV is a “profoundly political challenge, because it necessitates challenging the unequal social, political, and economic power of men and women and the ways in which this inequality is perpetuated through institutions at all levels of society” (Ward, 2005, p. 6).

THE IMPORTANCE OF GBV AS A YOUTH REPRODUCTIVE HEALTH ISSUE

- **Youth at Risk.** Rates of physical violence by male intimate partners against young women are also high. A 2008 analytical study conducted by the Demographic Health Survey (DHS) pertaining to intimate partner violence among couples in 10 DHS countries found that women from nine of the 10 countries reported having forced sexual experiences. The data also show that a major factor related to experiences of sexual violence is age at first marriage. In countries such as Bangladesh, Bolivia, Dominican Republic, Kenya, Rwanda, and Zimbabwe, women who were younger than 20 years old when they first married or started living with their current husband or partner were more likely to report physical or sexual violence than those who reported being 20 years or older when they first married (Hindin et al., 2008). According to a World Health Organization study (WHO, 2005), 20–36 percent of women ages 15–19 years old in 10 of 15 study sites reported being subject to at least one act of physical violence in the previous year. Moreover, rates of current physical violence are higher in the 15–19 year age group, compared with older women ages 20–49 years old.
- **Common Forms of GBV Particular to Youth.** The nature and extent of specific types of GBV can vary across cultures, countries, and regions. Some of the more common forms that involve young people include sexual exploitation, forced sex with children, early marriage, female genital mutilation/cutting, human trafficking, cross-

generational sex, and GBV in schools (Betron and Doggett, 2006).¹ Documentation and data on the following three forms of GBV show clear links with reproductive health outcomes for youth:

- **Female Genital Cutting/Mutilation.** “An estimated 100 million to 140 million girls and women worldwide have undergone female genital mutilation/cutting (FGM/C)” (WHO, 2008a, p. 1). Another estimate states that more than 3 million girls are at risk for genital cutting each year on the African continent alone” (Yoder and Khan, 2007). “Female Genital Cutting (FGC) is the name given to traditional practices that involve the partial or total cutting away of the female external genitalia and/or other injury to the female genitals, whether for cultural or non-therapeutic reasons” (Feldman-Jacobs and Clifton, 2008; USAID, 2004). FGC is customarily practiced on girls between the ages of 4 and 12 years old, albeit in some cultures, it can be performed as early as a few days after birth or as late as just prior to marriage, during pregnancy, or after the first birth (PRB, 2005). Immediate health consequences of FGC at any age can include severe pain, shock, hemorrhage, tetanus or sepsis, urine retention, open sores in the genital region, and injury to nearby genital tissue. A 2006 WHO study found an association with FGC and increased complications in childbirth and even maternal deaths. Specific side effects of childbirth include severe pain, hemorrhage, tetanus, infection, infertility, cysts and abscesses, urinary incontinence, and psychological and sexual problems (WHO, 2008a).
- **Early Marriage.** “Early marriage is the marriage of a person at an age at which she/he is not fully able to consent to the marriage and/or marriage at an age which results in vulnerability to reproductive health problems, psychosocial damage, or denial of education.” (Betron and Doggett, 2006, p. 10).” Societal norms, social status, isolation, and gender dynamics related to early marriage inhibit women and girls from having control over their own bodies and sexual and reproductive decisions. This can result in severe reproductive health problems, such as sexually transmitted infections; HIV; early childbearing (when the mother’s age is below the age of 18), leading to increased risk of maternal and infant mortality; unintended pregnancies; obstetric fistula; and other physical, psychological, and emotional damage associated with early childbearing (McIntyre, 2006; WHO and UNAIDS, 2008).

GBV-related Health Consequences for Youth

GBV results in reproductive health problems—often with lasting physical, social, emotional, psychological, and economic consequences (IGWG/USAID, 2008; Patel et al., 2007; UN General Assembly, 2006; WHO, 2005). Research has uncovered numerous links between exposure to GBV and subsequent health outcomes among youth:

Fatal Outcomes	Nonfatal Outcomes		
	Physical	Sexual and Reproductive	Psychological and Behavioral
<ul style="list-style-type: none"> ▪ Femicide ▪ Suicide ▪ AIDS-related mortality ▪ Maternal mortality 	<ul style="list-style-type: none"> ▪ Fractures ▪ Chronic pain syndromes ▪ Fibromyalgia ▪ Permanent disability ▪ Gastro-intestinal disorders 	<ul style="list-style-type: none"> ▪ Sexually transmitted infections, including HIV ▪ Unintended pregnancy ▪ Pregnancy complications ▪ Traumatic gynecologic fistula ▪ Abortion complications 	<ul style="list-style-type: none"> ▪ Depression and anxiety ▪ Eating and sleep disorders ▪ Drug and alcohol abuse ▪ Poor self-esteem ▪ Post-traumatic stress disorder ▪ Self-harm

Note: Adapted from Heise et al., 1999.

¹ Here, a child is defined per the UN Convention on the Rights of the Child as a person under the age of 18. The terms “young person” and “youth” lack similar legal definitions but here refer to adolescents and young adults.

- **Cross-generational Sex.** Cross-generational sex is often known as the “sugar daddy syndrome,” in which a young woman age 15-19 years old has a sexual relationship with a partner 5-10 years older in exchange for money or goods (Feldman-Jacobs and Worley, 2008). A 2008 DHS analytical study (Khan and Mishra, 2008) found that 12-29 percent of adolescent girls between the ages of 15-19 years old from Chad, Nigeria, and Senegal reported having high-risk sex in the past 12 months with a man 10 years or more older. In Uganda, the risk of HIV infection doubles for girls ages 15–19 years old who engage in sexual relations with male partners 10 or more years older (IWHC, 2008). Consequences of an unequal balance in age, gender, economic status, self-esteem, and power are believed to hinder adolescent girls’ ability to negotiate safe sexual practices. Cross-generational sex is not limited to sub-Saharan Africa, but most research on the practice has been conducted in that region. Studies from several countries in sub-Saharan Africa found that girls’ motivations for engaging in sexual relationships with older partners range from the need to find love, a spouse, and/or economic security; pay for school fees and supplies; and/or elevate their status among peers through material goods. Moreover, these girls may not realize the risks and severe health consequences (e.g., HIV, sexually transmitted infections, violence, unintended pregnancy, and other dangers) associated with these relationships (Hope, 2007).

KEY AREAS FOR POLICY ACTION

Effective action involves addressing both the complex root causes of GBV, as well as its immediate and long-term effects on victims. Reducing GBV requires countries to enact multisectoral policies, using myriad legal and policy instruments. Aside from the health sector, the education sector can play an important role in preventing and addressing GBV, particularly through health programs and policies in schools (Bott et al., 2005; Mirsky, 2005). In addition, the judicial sector and police can also play important roles by enforcing laws and policies aimed at GBV prevention and treatment. A comprehensive approach to addressing GBV as a youth reproductive health issue should include the following [integrated and multisectoral](#) policy actions:

- **Enact Policies to Empower Women and Girls.** To reduce gender imbalances at the root of GBV, key legal and policy actions should advocate for delayed marriage (IIPS and Population Council, 2008), promote equitable divorce and property laws that allow women the chance to leave abusive relationships (Global AIDS Alliance, 2007), and establish constitutional frameworks guaranteeing substantive equality for women (UN General Assembly, 2006a).
- **Pass Laws to Make Sexual Coercion and Domestic Violence Illegal.** It is just as important to have support for a legal and justice system that enforces these laws effectively (UN General Assembly, 2006a).
- **Demonstrate Clear Political Commitment to Ending GBV.** High-level government officials should consistently and publicly denounce GBV and support necessary changes in community norms that influence GBV-related behaviors of boys and young men (UN General Assembly, 2006b).
- **Establish Operational Policies and Guidelines to Support Program Efforts.** Public and private health facilities should institute policies and procedures to help providers recognize the signs of GBV and respond appropriately to meet the needs of GBV victims (IGWG/USAID, 2008).
- **Ensure that National Reproductive Health, HIV, Adolescent Health, and Maternal Health Policies and Legislation Specifically Address the Negative Reproductive Health Consequences of GBV.** Such policies should give clear guidance to health workers on their obligations in reporting and treating GBV (IGWG/USAID, 2008). Policies should also address the provision of [emergency contraception](#) and post-exposure antiretroviral prophylaxis to rape victims to protect against pregnancy and HIV infection.
- **Ensure that School Policies and Guidelines Directly Address GBV.** Schools may be a particularly unsafe place for young women (see box). Policies should strengthen the ability of teachers and administrators to address GBV (Mirsky, 2005) and also require them to report sexual violence against students (IGWG/USAID, 2008; USAID, 2007).

THE STATE OF POLICYMAKING

Although many governments have adopted policies related to GBV, few of them specifically relate to young people. Furthermore, most supportive laws and policies are not enforced (Bott et al., 2005; Harvard SPH, 2006).

Examples of national policies that refer to GBV and youth include Kenya's 2003 [Adolescent Health and Development Policy](#), which addresses the reporting of adolescent rape and punishment of offenders. The [2003 National AIDS Policy in Malawi](#) deals with sexual abuse of girls and women. [Panama's 1998 Framework for Formulation of Youth Policy and Action Plans](#) has a section on sexual abuse of youth and intra-family physical violence against adolescents.

Education Policies. South Africa is one of the few countries with a concerted policy effort to address GBV in schools (Bott et al., 2005). Its [1999 education sector policy on HIV/AIDS](#) promotes education to help students "protect themselves against rape, violence, inappropriate sexual behavior, and contracting HIV" (South Africa, 1999: p. 4) and mandates that the HIV/AIDS curriculum emphasize "the role of drugs, sexual abuse and violence, and sexually transmitted diseases in the transmission of HIV, and empowering learners to deal with these situations" (South Africa, 1999: 6–14). South Africa's 2007 [National Policy for the Prevention and Management of Learner Pregnancy](#) acknowledges the role and importance of teachers in educating students about sexual abuse, rape, and GBV and also directs school administrators to report cases of rape to the police.

Health Service Policies. Some organizations have adopted specific operational policies and procedures to address GBV among youth in clinical settings. Health nongovernmental organizations in the Dominican Republic, Peru, and Venezuela trained staff and instituted new clinical history forms, policies and procedures, and in-house services and referrals (Goldberg, 2006). The very first national policy for youth and sexual reproductive health was developed by Nicaragua. Developed by the government in 1996, [The National Integrated Child and Adolescent Policy](#) integrates reproductive health into a broader framework of citizenship, peer education, and political participation. Following a nationwide consultation with adolescents, the government integrated adolescent reproductive health needs into its poverty reduction strategy. In 2002, Nicaragua also developed a [National Policy against the Sexual Exploitation of Children and Adolescents](#).

Search for more GBV-related policies in the [Youth-Policy Database](#).

RESEARCH FINDINGS TO SUPPORT POLICY DEVELOPMENT

- [Guidelines for Integrating Sexual and Reproductive Health into the HIV/AIDS Component of Country-Coordinated Proposals to Be Submitted to the Global Fund to Fight AIDS, Tuberculosis, and Malaria, Round 7 and Beyond](#) (Doupe, 2007). Prepared by the Global AIDS Alliance and partners.
- *Intimate Partner Violence among Couples in 10 DHS Countries: Predictors and Health Outcomes* (Hindin et al., 2008). This report analyzes DHS data from 10 countries: Bangladesh, (2004), Bolivia (2003-2004), Dominican Republic (2002), Haiti (2005), Kenya (2003), Malawi (2004), Moldova (2005), Rwanda (2005), Zambia (2001-2002), and Zimbabwe (2005-2006). The report focuses mainly on health outcomes in relation to intimate partner violence in married and cohabiting relationships.
- [Journal of Adolescent Health, special issue focusing on sexual violence](#), various authors (JAH 2005), subscription only. This edition has several articles that highlight factors associated with rape and coercive sex, including sexually transmitted infections, other forms of violence, early menarche, and psychosocial problems.
- [Multi-Country Study on Women's Health and Domestic Violence against Women](#) (WHO, 2005). This study included interviews with 24,000 women in 15 sites in 11 countries (Bangladesh, Brazil, Ethiopia, Japan, Montenegro, Namibia, Peru, Samoa, Serbia, Tanzania, and Thailand). Report findings document the prevalence of intimate partner violence and its association with women's physical and mental health. The data cover nonpartner violence, sexual abuse during childhood, forced first sexual experience, women's responses to violence, and incidence of violence by age group.

- [MEASURE DHS Youth Corner](#). This website contains DHS reports that focus on data on various health factors related to youth sexual and reproductive health in 38 countries. The reports include information on issues that pertain to GBV (i.e., information on the proportion of girls and young women who experience physical violence, intimate partner violence, and nonconsensual sex).
- [National Cross-Sectional Study of Views on Sexual Violence and Risk of HIV Infection and AIDS among South African School Pupils](#) (Andersson et al., 2004). This survey of more than 200,000 South African students aged 10–19 years old found that both males and females accepted the reality of sexual coercion and had adopted attitudes to survive in a violent society.
- [Profiling Domestic Violence: A Multi-Country Study](#) (Kishor and Johnson, 2004). This report synthesizes DHS data from nine countries: Cambodia, Colombia, Dominican Republic, Egypt, Haiti, India, Nicaragua, Peru, and Zambia.
- [Female Genital Cutting in the Demographic and Health Surveys: A Critical and Comparative Analysis, DHS Comparative Reports](#) (Yoder et al., 2004). This comparative report summarizes DHS country data on FGC that were collected between 1989 and 2002. It gives the reader access to basic information on the distribution and practice of FGC.

FROM RHETORIC TO REALITY: POLICY IMPLEMENTATION TIPS

At the national level, many governments fail to formulate and implement effectively policies that express their commitment in addressing GBV (Bott et al., 2005). The Africa’s Health in 2010 project conducted a desk review about domestic violence based on DHS data in seven sub-Saharan African countries. The review sought to answer the following questions: “1. Is GBV addressed in the national planning documents (e.g., National Plans of Action [NPAs] and first and second generation Poverty Reduction Strategy Papers [PRSPs])? 2. In what sections is GBV addressed in the national planning documents, and is it included in the list of priorities? 3. Have DHS findings on domestic violence informed policies and programs in the national planning documents?” (Borwankar et al., 2008, p. 8).

The results of the review found the following:

- Five of the six country PRSPs reviewed mentioned GBV, but only two countries addressed GBV in a detailed manner. Of the four NPAs reviewed, Uganda and Zambia’s have a more complete approach to GBV compared with the PRSPs.
- GBV is not considered a priority in national planning documents and is usually mentioned in the narrative or under the cross-cutting gender and security law enforcement sections. Furthermore, “little emphasis is placed on addressing GBV as a public health issue (including to the linkages with HIV/AIDS, reproductive health) and to school-related GBV” (Borwankar et al., 2008, p. 11). Some countries cite DHS data and other health indicators.
- However, the data from the DHS domestic violence module were not cited in any of the national planning documents.

The review concluded that, despite the evidence cited on GBV prevalence in these countries, it is still not seen as an issue that merits national-level priority. The review stressed that it is imperative for governments and national stakeholders to recognize the importance of GBV as a health-related concern and allocate resources to a national GBV response that is multisectoral (sectors should include HIV/AIDS, reproductive health, mental health, education, law, criminal justice, human rights, social welfare, and gender) at the policy and program levels.

WATCH OUT FOR...

Well-intentioned GBV-related Policies that Could Negatively Affect Provision of Reproductive Health Services. Laws requiring health workers to report GBV cases can run counter to principles of confidentiality. Without clear guidelines and careful training of health workers, such policies may inadvertently diminish the willingness of both victims and health workers to discuss violence (Bott et al., 2005). Health workers must be trained to recognize the symptoms of GBV and assist victims of GBV in a knowledgeable manner. Victims should feel secure that health workers will not blame them or divulge confidential information to relatives, thus putting the victims at additional risk of violence. Furthermore,

HIV prevention services, sexual and reproductive health services, antenatal care, maternal and child health services, and voluntary counseling and testing services need to address GBV in an integrated manner (Doupe, 2007).

FAQs

Q. Can boys and men also be the victims of gender-based violence?

Yes. Traditional gender norms also fuel much violence that is experienced by men. This type of violence should also be recognized as a form of gender-based violence. According to WHO, “Rape and other forms of sexual coercion directed against men and boys take place in a variety of settings, including in the home, the workplace, schools, on the streets, in the military and during war, as well as in prisons and police custody” (Betron et al., 2006, p. 23). GBV against men tends particularly to occur in situations where men do not live up to “masculine” standards. For example, men who have sex with men who are outwardly open about their homosexuality have suffered violence and social ostracism. In Kenya, nearly 40 percent of men who have sex with men reported having been raped outside their home and 13 percent report having been assaulted by the police (Barker and Ricardo, 2005).

Q. Which international legal instruments address gender-based violence?

Several international agreements and policies address GBV and can be useful for formulating national laws and policies and conducting advocacy:

- [Convention on the Elimination of All Forms of Discrimination against Women](#) (CEDAW) (1979)
- [Convention on the Rights of the Child](#) (1990)
- The World Conference on Human Rights [Vienna Declaration and Programme of Action](#) (1993)
- [The UN Declaration on the Elimination of Violence against Women](#) (1993)
- [International Conference on Population and Development Programme of Action](#) (1994)
- [Inter-American Convention on the Prevention, Punishment, and Eradication of Violence against Women](#) (1994)
- UN Fourth World Conference on Women [Beijing Declaration](#) and [Platform for Action](#) (1995)
- Organization of African Unity [Protocol to the African Charter on Human and People’s Rights on the Rights of Women in Africa](#) (1995)

For more on international policies, see [Guedes, 2004](#); [Bott et al., 2005](#); [Prevent GBV Africa](#) website; and [UN Division for the Advancement of Women](#) website.

RELATED LINKS AND RESOURCES

Documents

- [Addressing Gender-Based Violence from the Reproductive Health/HIV Sector: A Literature Review and Analysis](#) (Guedes, 2004). This USAID-commissioned literature review analyzes programs in developing countries and includes chapters on policy and on youth-focused programming.
- [Addressing Gender-Based Violence through USAID’s Health Programs: A Guide for Health Sector Program Officers](#) (IGWG/USAID, 2008). This guide is intended to help USAID program officers integrate GBV initiatives into their health sector portfolios during project design, implementation, and evaluation. The guide includes sections on youth and policy.
- [In-Depth Study on All Forms of Violence against Women: Report of the Secretary-General](#) (UN General Assembly, 2006a). This study summarizes laws and policies that are effective in combating violence against women.
- [Preventing and Responding to Gender-Based Violence in Middle- and Low-Income Countries: A Global Review and Analysis](#) (Bott et al., 2005). This World Bank Policy Research Working Paper includes detailed analysis of policy and legal reform.

- [Programming to Address Violence against Women: 10 Case Studies](#) (UNFPA, 2007). This volume documents UNFPA’s experience addressing violence against women in Bangladesh, Colombia, Ghana, Kenya, Mauritania, Mexico, Morocco, Romania, Sierra Leone, and Turkey. The volume includes descriptions of policy initiatives and programs serving adolescents.
- [Responding to Gender-Based Violence: A Focus on Policy Change](#) (POLICY Project, 2006). This toolkit is a resource for advocates to raise awareness and influence policy.
- [Sex without Consent: Young People in Developing Countries](#) (Jejeebhoy et al., 2005, abstract only). This book documents the extent of nonconsensual sex among youth, key factors that put youth at risk, evidence-based directions for programming, and legal and policy responses.
- “The Vulnerability of Adolescence: Legal Responses to Non-Consensual Sexual Experiences of Young Persons in India” (Jaising, 2005).
- [Young Men and Masculinity in Sub-Saharan Africa: Implications for HIV/AIDS, Conflict, and Violence](#) (Barker and Ricardo, 2005). This document describes programs that apply a gender perspective to young men and attempt to change norms around negative male behaviors, including GBV. Some of these programs also focus on policy change.

Websites

- [Collation of Laws and Resources on Domestic Violence](#). Maintained by the Asia-Pacific Forum on Women, Law, and Development and intended as a resource for groups and communities undertaking legal reform on domestic violence.
- [End Violence against Women](#). Information clearinghouse maintained by the USAID | INFO Project.
- [Gender-Based Violence Prevention Network](#). Focused on East and Southern Africa and includes a [section on youth](#).
- [Interagency Gender Working Group \(IGWG\)](#). This USAID-supported group promotes gender equity within population, health, and nutrition programs worldwide. Website includes a [section on youth](#).
- [Interagency Youth Working Group \(IYWG\)](#). Includes a searchable resource database with links to more than 40 documents relating to nonconsensual sex.
- [Sexual Violence Research Institute \(SVRI\)](#). This important website contributes to eliminating sexual violence by addressing the lack of research on the different aspects of sexual violence, as well as drawing attention to including policymakers and the media to this issue.
- [United Nations Division for the Advancement of Women, Violence against Women Web Page](#). Contains links to several policy documents.

REFERENCES

- African Child Policy Forum. 2006. [Born to High Risk: Violence against Girls in Africa](#). A report prepared for the Second International Policy Conference on the African Child: Violence against Girls in Africa, May 11–12, 2006. Addis Ababa, Ethiopia.
- Andersson, N., A. Ho-Foster, and J. Mathis. 2004. National Cross-sectional Study of Views on Sexual Violence and Risk of HIV Infection and AIDS among South African School Pupils. *British Medical Journal* 2004; 329:952 (23 October).
- Barker, G., and C. Ricardo. 2005. “Young Men and the Construction of Masculinity in Sub-Saharan Africa: Implications for HIV/AIDS, Conflict, and Violence.” *Social Development Papers, Conflict Prevention and Reconstruction* No. 26. Washington, DC: World Bank.
- Betron, M., and E. Doggett. 2006. *Linking Gender-Based Violence Research to Practice in East, Central, and Southern Africa: A Review of Risk Factors and Promising Interventions*. Washington, DC: POLICY Project, Futures Group.
- Borwankar R., R. Diallo, and A. E. Sommerfelt. 2008. *Gender-Based Violence in Sub-Saharan Africa: A Review of Demographic and Health Survey Findings and Their Use in National Planning*. Washington DC: USAID/AFR/SD and Africa’s Health in 2010/AED.

- Bott, S., A. Morrison, and M. Ellsberg. 2005. "[Preventing and Responding to Gender-Based Violence in Middle- and Low-Income Countries: A Global Review and Analysis.](#)" *World Bank Policy Research Working Paper* 3618. Washington, DC.
- Doupe, Andrew. 2007. [Guidelines for Integrating Sexual and Reproductive Health into the HIV/AIDS Component of Country Coordinated Proposals to Be Submitted to the Global Fund to Fight AIDS, Tuberculosis, and Malaria Round 7 and Beyond.](#) Washington, DC: Global AIDS Alliance. Available at: http://aidsalliance.3cdn.net/ebe8657456c6f186b3_7tm6iio1b.pdf (accessed October 14, 2009).
- Feldman-Jacobs, C., and D. Clifton. 2008. *Female Genital Mutilation/Cutting: Data and Trends.* Washington, DC: Population Reference Bureau.
- Feldman-Jacobs, C., and H. Worley, 2008. *Cross-Generational Sex: Risks and Opportunities.* Washington, DC: Population Reference Bureau.
- Global AIDS Alliance. 2007. "Call to Action: Violence Free Zone: Ending School-Related Violence, Prevent HIV/AIDS." Washington DC: Global AIDS Alliance.
- Goldberg, Rachel. 2006. "[Living Up to Their Name: Profamilia Takes on Gender-Based Violence.](#)" *Quality/Calidad/Calite* No. 18: 5–31. New York: Population Council.
- Guedes, Alessandra. 2004. [Addressing Gender-Based Violence from the Reproductive Health/HIV Sector: A Literature Review and Analysis.](#) Washington, DC: The Population Technical Assistance Project/ Inter-Agency Gender Working Group.
- Harvard School of Public Health (SPH). 2006. "HIV/AIDS and Gender-Based Violence (GBV) Literature Review." Boston, Massachusetts. Available at: http://www.hsph.harvard.edu/pihhr/files/Final_Literature_Review.pdf (accessed September 8, 2009).
- Heise, L., M. Ellsberg, and M. Gottemoeller. 1999. "Ending Violence against Women." *Population Reports* XXVII (Series L, Number 11). Available at: www.inforforhealth.org/pr/111/violence.pdf (accessed October 14, 2009).
- Hindin, M., S. Kishor, and D. Asara. 2008. "Intimate Partner Violence among Couples in 10 DHS Countries: Predictors and Health Outcomes." *DHS Analytical Studies* No. 18. Calverton, MD: Macro International.
- Hoffman, S., L. O'Sullivan, A. Harrison, C. Dolezal, and A. Monroe-Wise. 2006. "[HIV Risk Behaviors and the Context of Sexual Coercion in Young Adults' Sexual Interactions: Results from a Diary Study in Rural South Africa.](#)" *Sexually Transmitted Diseases* 33(1): 52–58.
- Hope, R. 2007. *Addressing Cross-Generational Sex: A Desk Review of Research and Programs.* Washington, DC: Population Reference Bureau.
- Inter-Agency Gender Working Group (IGWG)/USAID. 2008. [Addressing Gender-Based Violence through USAID's Health Programs: A Guide for Health Sector Program Officers.](#) Washington, DC: IGWG.
- International Institute for Population Sciences (IIPS) and the Population Council. 2008. "[How Early Marriage Compromises Girls' Lives, Maharashtra.](#)" *Youth in India: Situation and Needs 2006–2007.* Policy Brief No. 6. Mumbai.
- International Women's Health Coalition (IWHC). 2008. *Triple Jeopardy: Female Adolescence, Sexual Violence and HIV/AIDS.* New York: IWHC.
- Jaising, Indira. 2005. "The Vulnerability of Adolescence: Legal Responses to Non-Consensual Sexual Experiences of Young Persons in India." Ch. 14, pp. 216-26. In Jejeebhoy et al. (2005).

Jejeebhoy, S., I. Shah, and S. Thapa, eds. 2005. [*Sex without Consent: Young People in Developing Countries*](#). New York: Zed Books.

[*Journal of Adolescent Health \(JAH\)*](#). 2005. [Special Issue Focusing on Sexual Violence](#) 36(5): 359–448.

Khan, S., and V. Mishra. 2008. “Youth Reproductive and Sexual Health.” *DHS Comparative Reports* No. 19. Calverton, MD: Macro International.

Kishor, S., and K. Johnson. 2004. [*Profiling Domestic Violence: A Multi-Country Study*](#). MEASURE DHS. Calverton, MD: ORC Macro.

McIntyre, P. 2006. *Married Adolescents: No Place of Safety*. Geneva: WHO.

Mirsky, J. 2005. “Developing Opportunities within the Education Sector to Prevent Non-Consensual Sexual Experiences: An Emerging Issue for Human Rights, Public Health, and Education Development Goals.” In Jejeebhoy et al. (2005).

Patel, V., A. Flisher, S. Hetrick, and P. McGorry. 2007. “Mental Health of Young People: A Global Public-Health Challenge.” *The Lancet* 369: 1302–13.

POLICY Project. 2006. *Responding to Gender-Based Violence: A Focus on Policy Change A Companion Guide*. Washington, DC: Futures Group, POLICY Project.

Population Reference Bureau (PRB). 2005. *Abandoning Female Genital Mutilation/Cutting: Information from around the World*. Washington, DC: PRB.

South Africa. 1999. “[National Policy on HIV and AIDS for Learners and Educators in Public Schools and Students and Educators in Further Education and Training Institutions.](#)” Vol. 410, No. 20372. Pretoria: Department of Education.

South Africa. 2007. [Measures for the Prevention and Management of Learner Pregnancy](#). Pretoria: Department of Education.

UN General Assembly. 2006a. [In-Depth Study on All Forms of Violence against Women: Report of the Secretary-General](#). New York: UN.

UN General Assembly. 2006b. [Rights of the Child: Report of the Independent Expert for the United Nations Study on Violence against Children](#). New York: UN.

United Nations Population Fund (UNFPA). 2007. [Programming to Address Violence against Women: 10 Case Studies](#). New York: UNFPA.

UNFPA. 1998. “Violence Against Girls and Women: A Public Health Priority.” UNFPA Gender Theme Group, Interactive Population Center. New York: UNFPA.

United Nations Program on HIV/AIDS (UNAIDS). 2006. “[Preventing HIV/AIDS in Young People. A Systematic Review of the Evidence from Developing Countries.](#)” *WHO Technical Report Series* 938. Inter-agency Task Team on Young People. Geneva: WHO.

United States Agency for International Development (USAID). 2004. “Bureau for Global Health Strategy for Female Genital Cutting (FGC), FY2004-2006.” Washington, DC: USAID.

USAID. 2007. *The Safe Schools Program: Student and Teacher Baseline Report on School-Related Gender-Based Violence in Machinga District, Malawi*. Washington, DC: DevTech Systems.

USAID. 2009. "A Guide to Programming Gender-Based Violence Prevention and Response Activities: Draft for Discussion Purposes." Washington, DC: Gender-Based Violence Working Group.

Ward, J. 2005. *Because Now Men Are Really Sitting on Our Heads and Pressing Us Down...*

Report of a Preliminary Assessment of Gender-Based Violence in Rumbek, Aweil (East and West), and Rashad County, Nuba Mountains. Supported by the USAID-USDA PASA in collaboration with the University of Missouri, Tuskegee University, and University of Maryland-Eastern Shore.

World Health Organization (WHO). 2005. [Multi-Country Study on Women's Health and Domestic Violence against Women](#). Geneva: WHO.

WHO. 2008a. *Eliminating Female Genital Mutilation: An Interagency Statement*. Geneva: WHO.

WHO. 2008b. "Female Genital Mutilation." *WHO Fact Sheet* No. 241. Geneva: WHO.

WHO and UNAIDS. 2008. *Technical Guidance for Global Fund HIV Proposals Cross-Cutting Issues: Addressing Gender Inequalities: Strengthening HIV/AIDS Programming for Women and Girls*. Geneva: WHO and UNAIDS.

Yoder, P. S., and S. Khan. 2007. *Numbers of Women Circumcised in Africa: The Production of a Total*. Calverton, MD: Macro International.

Yoder, P. S., N. Abderrahim, and A. Zhuzhuni. 2004. "Female Genital Cutting in the Demographic and Health Surveys: A Critical and Comparative Analysis. DHS Comparative Reports." *DHS Comparative Reports* No. 7. MEASURE DHS. Calverton, MD: ORC Macro.

For More Information

Health Policy Initiative, Task Order I
Futures Group
One Thomas Circle, NW, Suite 200
Washington, DC 20005 USA
Tel: (202) 775-9680
Fax: (202) 775-9694

policyinfo@futuresgroup.com
<http://www.healthpolicyinitiative.com>
<http://ghiqc.usaid.gov>

The USAID | Health Policy Initiative, Task Order I, is funded by USAID under Contract No. GPO-I-01-05-00040-00, beginning September 30, 2005. HIV-related activities are supported by the President's Emergency Plan for AIDS Relief. Task Order I is implemented by Futures Group, in collaboration with the Centre for Development and Population Activities (CEDPA), White Ribbon Alliance for Safe Motherhood (WRA), and Futures Institute.

The views expressed in this publication do not necessarily reflect the views of USAID or the U.S. government.